

Gauss Elimination

Zvi Rosen
Department of Mathematics

October 11, 2016

Guiding Question

Given a matrix equation $Ax = b$, for a real matrix A and a real vector b , how do we find a solution x ?

Graphical Method

Consider each equation as defining a line / plane / hyperplane in Euclidean space.

Draw each plane and see where they all intersect.

Cramer's Rule

For $Ax = b$, let $A_j =$ the matrix formed by replacing the j -th column of A with b . Then:

$$x_j = \frac{\det(A_j)}{\det(A)}.$$

What is the determinant of a matrix? See next slide.

Determinants


The determinant of a square matrix A can be defined inductively as follows:

1. The determinant of a 1×1 matrix is its only entry.
2. The determinant of a $n \times n$ matrix can be obtained by: Fixing a row j . Defining the matrix $A(j, k)$ to be the matrix leaving out the j -th row and k -th column.

$$\det(A) = \sum_{k=1}^n (-1)^{j+k} A_{j,k} \det(A(j, k))$$

Determinants – Geometric View

If you consider the matrix A as a linear map between two vector spaces, then the determinant can be thought of as the volume of the unit cube after passing through the map A .



–image from Wikipedia.

Naive Gauss Elimination

INPUT: $n \times (n + 1)$ Matrix $A|b$

1. Fix pivot variable $A_{1,1}$.
2. Perform **row operations** to eliminate everything below the pivot variable.
3. If $k < n$, change pivot variable from $A_{k,k}$ to $A_{k+1,k+1}$ and repeat step 2.
4. Solve for x_k and substitute its value into Row $k - 1$.
Repeat to solve for each x_k .

Row operations: Permute rows, add a scalar multiple of row j to row k .

Naive Gaussian Elimination - Example

$$\left(\begin{array}{ccc|c} 1 & 2 & 3 & 4 \\ 6 & 5 & 8 & 7 \\ 3 & 1 & 4 & 2 \end{array} \right)$$

Naive Gauss - Complexity

How many operations does Naive Gaussian elimination take?

$$\begin{aligned}\# \text{operations} &= \sum_{k=1}^{n-1} \#(\text{operations at pivot } k) \\ &+ \sum_{k=1}^n \#(\text{operations to solve for } x_k) \\ &= \sum_{k=1}^{n-1} 2(n-k)(n-k-1) + \sum_{k=1}^n (n-k+2)\end{aligned}$$

Naive Gauss - Complexity

$$\begin{aligned} &= \sum_{k=1}^{n-1} 2n^2 - \sum_{k=1}^{n-1} 4nk + \sum_{k=1}^{n-1} 2k^2 - \sum_{k=1}^{n-1} 2n \\ &+ \sum_{k=1}^{n-1} 2k + \sum_{k=1}^n n - \sum_{k=1}^n k + \sum_{k=1}^n 2. \\ &= 2n^3 - 3n^2 + 3n + (-4n + 1) \sum_{k=1}^{n-1} k + 2 \sum_{k=1}^{n-1} k^2. \end{aligned}$$

Note that $\sum_{k=1}^{n-1} k = \frac{1}{2}n^2 + O(n)$ and $\sum_{k=1}^{n-1} k^2 = \frac{1}{3}n^3 + O(n^2)$.

Naive Gauss - Complexity

So we are left with:

$$\begin{aligned} &= 2n^3 - 3n^2 + 3n + (-4n + 1)\left(\frac{1}{2}n^2 + O(n)\right) + 2\left(\frac{1}{3}n^3 + O(n^2)\right) \\ &= \frac{2}{3}n^3 + O(n^2). \end{aligned}$$

Pivoting

We have to be careful! Sometimes $A_{k,k} = 0$ or is numerically very close to zero.

- ▶ Partial pivoting means at each step, search column k for the largest element A_{jk} then switch rows j and k .
- ▶ Complete pivoting means that we also search for the highest value in the row j (This would mean relabeling variables, which is undesirable).

Pivoting - Example

Example

$$\begin{pmatrix} .0003 & 3.0000 \\ 1.0000 & 1.0000 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 2.0001 \\ 1.0000 \end{pmatrix}$$

Compare the results you obtain via naive Gaussian elimination, and pivoting. (Use MATLAB for your computation.)

Tridiagonal Systems

A tridiagonal system is a matrix equation $Ax = b$, when the only nonzero entries of A are $A_{i,i-1}$, $A_{i,i}$, and $A_{i,i+1}$ (wherever these indices make sense).

$$\begin{pmatrix} a_{1,1} & a_{1,2} & 0 & 0 & \cdots & 0 \\ a_{2,1} & a_{2,2} & a_{2,3} & 0 & \cdots & 0 \\ 0 & \ddots & \ddots & \ddots & & \vdots \\ 0 & \cdots & \cdots & 0 & a_{n-1,n} & a_{n,n} \end{pmatrix}$$

Tridiagonal Systems - Complexity

Question: How many operations does Gaussian Elimination take for a tridiagonal system?

Perform a complexity computation similar to the slides above regarding Naive Gaussian Elimination, accounting for both elimination and substitution steps. Your result should be $O(n)$.

LU Factorization

Problem: Find a way to quickly solve $Ax = b$ when A is fixed but we want to solve for a number of inputs b .

Idea: Factor A as $A = LU$ where L is lower-triangular and U is upper-triangular. Then, suppose there exists a vector d so that $Ld = b$.

$$Ax = b \implies L U x = b \implies L U x = L d \iff U x = d.$$

At this point, we have two equations with triangular matrices:

$$Ux = d, Ld = b.$$

Solving with LU Factorization

Given these two triangular systems $Ux = d$, $Ld = b$, we do not need to worry about the elimination part of Gaussian elimination – just the substitution part.

Solve for d_1 through d_n substituting down at each step.
Then solve for x_n through x_1 substituting up at each step.
This drastically cuts down complexity.

LU Factorization – How?

It should be clear by now that an LU factorization would be really useful for equation solving. But how do we obtain it? Fix A as our matrix.

Perform the standard Gaussian elimination, when our pivot variable is A_{kk} , we eliminate entry A_{jk} below the diagonal by replacing row R_j with $c_{jk} * R_k + R_j$. Set L_{jk} to be $-c_{jk}$.

- ▶ U is the matrix spit out by Gaussian elimination.
- ▶ L is the matrix with ones on the diagonal, and the lower entries obtained as above.

LU Factorization - Example

Let A be the matrix:

$$\begin{pmatrix} 1 & 2 & 3 \\ 6 & 5 & 8 \\ 3 & 1 & 4 \end{pmatrix}$$

Let b be the vector $(4, 7, 2)$. Solve $Ax = b$ by:

1. Standard Gaussian Elimination.
2. Finding $A = LU$ then solving $Ld = b$ and $Ux = d$.