

Quiz 1

MATH 1B, SPRING 2012

GSI: Zvi Rosen

SECTION:

NAME: SOLUTION

Solve the following integral, showing all steps clearly:

$$\int_0^a x^2 \sqrt{a^2 - x^2} dx$$

Let $x = a \sin \theta$, where $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$.

Then, $dx = a \cos \theta d\theta$.

To change the bounds, $x = a \sin \theta = a$ when $\sin \theta = 1$,

i.e. when $\theta = \frac{\pi}{2}$. $x = a \sin \theta = 0$ when $\sin \theta = 0$, at $\theta = 0$.

So, our integral becomes:

$$\begin{aligned} & \int_0^{\pi/2} a^2 \sin^2 \theta \sqrt{a^2(1-\sin^2 \theta)} \cdot a \cos \theta d\theta. && \text{use } 1-\sin^2 \theta = \cos^2 \theta \\ & = \int_0^{\pi/2} a^4 \sin^2 \theta \cos \theta |\cos \theta| d\theta. && \left. \begin{array}{l} \bullet \text{ On this domain, } \cos \theta > 0, \\ \text{so } |\cos \theta| = \cos \theta. \\ \bullet 2 \sin \theta \cos \theta = \sin 2\theta \\ \text{so, } \sin^2 \theta \cos^2 \theta = \frac{1}{4} \sin^2 2\theta \\ \bullet \sin^2 2\theta = \frac{1}{2}(1 - \cos 4\theta) \end{array} \right\} \\ & = \int_0^{\pi/2} a^4 \sin^2 \theta \cos^2 \theta d\theta. \\ & = \frac{a^4}{4} \int_0^{\pi/2} \sin^2 2\theta d\theta. \\ & = \frac{a^4}{4} \int_0^{\pi/2} \frac{1}{2}(1 - \cos 4\theta) d\theta \\ & = \frac{a^4}{8} \left[\theta - \frac{1}{4} \sin 4\theta \right]_0^{\pi/2} = \frac{a^4}{8} \cdot \left(\frac{\pi}{2} - 0 \right) = \frac{\pi a^4}{16}. \end{aligned}$$