

(Based on a worksheet by Rob Bayer)

### Second Order Equations

1. For each of the following pairs of functions, determine whether they are linearly independent or dependent:

(a)  $f(x) = e^x$ ,  $g(x) = e^{x+1}$

(c)  $f(x) = \ln(x^3)$ ,  $g(x) = \ln(x^\pi)$

(b)  $f(x) = xe^{2x}$ ,  $g(x) = e^{2x}$

(d)  $f(x) = \sin(x)$ ,  $g(x) = \cos(x)$

2. Which of the following second order differential equations are linear? Homogeneous?

(a)  $e^x y'' + \cos(3x^2)y' + 3y = 0$

(b)  $y'' + 3y' + 7y = \cos x$

(c)  $y'' + 3xy' + y^2 = 0$

(d)  $\tan(y'') + \cos(x)y' = e^x$

3. Consider the differential equation  $y'' = -y$

(a) By just thinking about it for a while, come up with two linearly independent solutions to this equation and then use them to find the general solution. (Think about elementary functions whose derivatives cycle.)

(b) Show that  $y = \cos(x + a)$  is a solution for any constant  $a$ .

(c) Argue that parts (a) and (b) show that  $\cos(x + a) = C_1 \cos x + C_2 \sin x$  for an appropriate choice of  $C_1$  and  $C_2$

(d) Without using trig identities, find  $C_1$  and  $C_2$ . (Hint: what should  $y(0)$  and  $y'(0)$  be? Remember that  $a$  is a constant, so  $C_1$  and  $C_2$  can refer to it.) Does this formula look familiar?

### Solving Second Order Homogeneous Linear ODEs with Constant Coefficients

1. Find the general solution to  $y'' + 3y' - 18y = 0$

2. Solve the initial value problem  $y'' + 4y' + 4y = 0$  with  $y(0) = 1$  and  $y'(0) = 3$

3. Solve the boundary value problem  $y'' = y$  with  $y(0) = 0$  and  $y(2) = 2$